




# Cell Biology




A cell is chemical system that is able to maintain its structure and reproduce. Cells are the fundamental unit of life. All living things are cells or composed of cells.


The interior contents of cells is the cytoplasm. The cytoplasm is isolated from the surrounding environment by the \_\_\_\_\_ . There are two fundamentally different forms of cells.

Prokaryotic cells - relatively simple cells - lack nuclear membrane and many organelles - bacteria and their relatives are all prokaryotic



Eukaryotic cells - more complex cells - have a nucleus and many organelles - all cells of plants, animals, fungi, and protists






Most cells are small

Prokaryotic: 1-10  $\mu\text{m}$


Eukaryotic: 10 - 100  $\mu\text{m}$

(1  $\mu\text{m} = .001 \text{ mm}$ )

Cells and organelles




# Organelles, Macromolecules, & Atoms



# Why are cells small?

As cell size increases the volume increases much faster than the surface area.


Cells obtain nutrients, gain information and rid waste through their plasma membrane.



As cell size increases, a cell's ability to exchange with its environment becomes limited by the amount of membrane area that is available for exchange.

Robert Hooke - 1665 -  
using an early microscope  
viewed cork and saw many  
repeating box-like  
structures and called them  
“cells.”

What he saw were spaces  
surrounded by walls that  
once contained living  
cells.



Since Hooke’s first observations  
what is known about cells has  
increased greatly.

# Cell Theory

- Cells are the fundamental unit of life - nothing less than a cell is alive.
- All organisms are constructed of and by cells.
- All cells arise from preexisting cells. Cells contain the information necessary for their own reproduction. No new cells are originating spontaneously on earth today.
- Cells are the functional units of life. All biochemical processes are carried out by cells.
- Groups of cells can be organized and function as multicellular organisms
- Cells of multicellular organisms can become specialized in form and function to carry out subprocesses of the multicellular organism.


# Prokaryotic cell structure

small, with a plasma membrane surrounded by a rigid **cell wall** -  
in many the cell wall is made of \_\_\_\_\_ - a carbohydrate  
cross-linked with polypeptides  
cell wall may be covered with a **capsule** made of polysaccharides

few or no membrane  
enclosed spaces  
within the cytoplasm

no nucleus - DNA is  
in a region called the  
**nucleoid**

DNA is circular and  
**naked** (has no protein  
associated with it)






Bacteria often have **flagella** with a single protein core (flagellin) that they can use to move in a rotary corkscrew like fashion


The rotary motor of prokaryotic flagella is powered by proton flow through the cell membrane.

Rotating structures are rare in nature.



Membrane enclosed spaces allow cell functions to be compartmentalized and isolated from other functions. Prokaryotes lack membrane enclosed spaces in their cytoplasm.

Some prokaryotes are photosynthetic. The biochemical machinery for trapping light energy is contained within a highly folded plasma membrane.



# Eukaryotic cell structure

larger, with a typical plasma membrane - some with a cell wall



Many \_\_\_\_\_ and other interior spaces enclosed by membranes:

**Nucleus, Endoplasmic reticulum, Golgi apparatus,**

**Mitochondria, Chloroplasts, Lysosomes, Vacuoles, Vesicles**

Cytoplasm with a cytoskeleton - protein tubules and fibers

cell wall found in plants (cellulose), fungi (chitin), some protists





# Cellular Organelles

**Nucleus** - the largest and most obvious membrane bound compartment - controls cell activities




contains the **nucleolus** - a darkened region where ribosomal RNA is synthesized

contains **chromosomes** - consist of DNA wrapped around proteins



Nucleus is surrounded by the nuclear envelope - a double membrane

Nuclear membrane has nuclear pores that control entry and exit of materials




## Chromosome - “colored body”


consists of both DNA and protein - seen as chromosomes when highly condensed in preparation for cell division

At other times the DNA and protein are threadlike and called \_\_\_\_\_.

The most common proteins are histones. DNA is coiled around histones in a regular pattern that produces structures called nucleosomes.




**Endoplasmic reticulum (ER)** - a web-like series of membranes within the cytoplasm in the form of flattened sheets, sacs, tubes, creates many membrane enclosed spaces - spreads throughout the cytoplasm - has connections with the outer membrane of the nucleus and the plasma membrane  
interior space is called the \_\_\_\_\_



**Functions:**

- circulation and transport
- storage of proteins and minerals
- synthesis of lipids, carbohydrates, and proteins
- A large surface area for enzyme action.

## Two types of ER - rough and smooth





**rough ER** - studded with ribosomes  
site of synthesis of many proteins  
all ribosomes on rER are actively involved in protein synthesis -

**smooth ER** - site for synthesis of steroids and other lipids  
Ca<sup>++</sup> storage in muscles  
detoxification of drugs, toxins, alcohol (especially in liver)

The highly convoluted surface provides a large surface area for enzymatic activities. Many enzymes are imbedded in the membranes.






## Ribosomes - protein

synthetic machinery

- two subunits - large and small - each made of protein and ribosomal RNA (rRNA)
- subunits associate when they are synthesizing proteins
- protein synthesis occurs on ribosomes that are free-floating in the cytoplasm and on ribosomes attached to ER
- rRNA is synthesized in the nucleolus

## Golgi Apparatus -

a collection of membranes associated with the ER  
composed of flattened sacs called \_\_\_\_\_  
concentrates and packages proteins synthesized on the ER




The Golgi is functionally associated with the ER.

Proteins synthesized on the ER are concentrated internally and transport vesicles are budded off

Transport vesicles fuse with the Golgi, dump their contents into the Golgi

Golgi packages proteins in vesicles so that they may be excreted from the cell, or used within the cell.




Secretory vesicles - used for excretion - leave the Golgi and move to plasma membrane where they fuse and dump their contents outside - seen in many glands

The Golgi Apparatus also forms lysosomes

Lysosomes - vesicles filled with digestive enzymes - used for intracellular digestion

Particles can be taken into cell by phagocytosis and vesicle fused with lysosome

The components of organelles can be recycled after digestion by lysosomes




## Microbodies: Peroxisomes and Glyoxisomes

vesicles that form through growth and division within the cytoplasm

**Glyoxisomes** are found in plants - contain enzymes that convert fats into carbohydrates

**Peroxisomes** - used for removing reactive compounds from the cytoplasm - create  $H_2O_2$  as a byproduct and degrade it with the enzyme catalase




**Mitochondria** - cellular powerhouses - the site of much of the energy harvest by cells

have double membrane structure

inner membrane folded into inward projections called **cristae**

two spaces within the mitochondrion - the **matrix** and the **intermembrane space**




## Mitochondria -

- The site of oxygen consumption within cells
- Have their own DNA that is similar to prokaryotic DNA
- Have their own ribosomes that are similar in construction to prokaryotic ribosomes
- Synthesize many, but not all, of their own proteins
- Mitochondria replicate by binary fission - similar to prokaryotic cell division

**Chloroplasts** - sites of photosynthesis - in nearly all plants and some protists  
trap light energy and convert it into chemical energy  
have double membrane structure - inner space is the stroma

Within the stroma have a series of stacks of flattened membrane structures called thylakoids - the stacks are called grana



The light energy trapping molecules of photosynthesis are found in the membranes of the thylakoids.



## Chloroplasts

have their own DNA, similar to prokaryotic DNA

Can synthesize many of their own proteins using prokaryote-like ribosomes

Synthesize many, but not all, of their own proteins

Replicate through division similar to prokaryotic cell division

## Chloroplasts can take on other functions

\_\_\_\_\_ synthesize and store starch in roots and tubers

\_\_\_\_\_ have pigments and give fruits ripened color


Centrioles - are part of specialized region of the cell called the centrosome (cell center)

found in animals and most protists  
the centrioles are involved in the production of microtubules

microtubules have many functions including moving chromosomes during cell division

centriole structure - 9 triplets of microtubules surrounding a hollow core - \_\_\_\_\_

similar to the basal body of flagella




Cytoskeleton - scaffolding of proteins that transport materials, position and move organelles, maintain and change cell shape, and organize enzymes into functional associations

3 components - actin filaments, microtubules, and intermediate filaments


all are polymers of smaller protein subunits -  
lengthen through addition  
of polymer subunits,  
shorten through

---



**actin filaments** - involved in cell movements and in membrane deformations - smallest components of the cytoskeleton

**microtubules** - hollow tubes made of proteins called **tubulins** responsible for cell movements and movements of organelles within the cytoplasm, movement of chromosomes during cell division - largest components of the cytoskeleton



**intermediate filaments**

- 8 stranded protein fibers - play a role in cell structure, anchoring organelles

and in transport of materials within the cytoplasm

anchor neighboring cells to each other in tissues

## Flagella and Cilia - cellular appendages

can propel cells or propel materials over the cell surface

cells that have flagella have few (usually 1 or 2)


cells that have cilia have many - covering the surface

flagella move with whip-like movements to propel the cell

cilia have a more regular stroke and groups of cilia appear to

move in unison, resulting in a wave-like motion

flagella 5 to 20x longer than cilia





## Structure

has basal body with 9 + 0 structure of microtubules

flagellum is membrane bound with pairs of microtubules in a 9+2 pattern

each pair of tubules has short arms of another protein - dynein - that extend to neighboring tubules


movement of the flagellum is produced by sliding of the microtubule pairs



Plant Cells have, in addition to the collection of organelles found in other groups, a central vacuole for storage and for producing \_\_\_\_\_ pressure inside the the cell.


The central vacuole is usually filled with water and solutes. A high solute concentration draws water into the vacuole, expanding the vacuole and the cell.

Because plant cells are enclosed by a cell wall, the expansion of the vacuole can exert pressure on the cell without causing the cell to burst.



Plants have cell walls made of cellulose.

During cell division plant cells build dividing walls between the two new cells called the **cell plate**. An adhesive layer - the **middle lamella** - is laid down between the new cell walls



Cell walls can be thickened through the addition of materials to the inside of the primary cell wall.




## Where did Eukaryotic Cells come from?

The oldest rocks with evidence of fossil cells date to 3.5 billion years. The oldest rocks with cells large and complex enough to be eukaryotic date to 1.0 billion years.


For 2.5 billion years only prokaryotic cells existed on earth.

The best hypothesis for the origin of eukaryotic cells was proposed by Lynn Margulis in the early 1970s. This hypothesis is now called the \_\_\_\_\_.

Eukaryotic cells appear to be the product of a collaboration among different types of prokaryotic cells. Some prokaryotic cells became the host for other prokaryotic cells that lived inside them. Some of the complex organelles of eukaryotes provide evidence for this theory.



Mitochondria and chloroplasts appear to be the direct descendants of energy producing bacteria. Mitochondria are the descendants of bacteria that were capable of oxidative respiration. Chloroplasts are the descendants of photosynthetic bacteria.



## Evidence:

Both have their own DNA and ribosomes that are similar to those found in prokaryotes. Both make many of their own proteins and both multiply in a fashion similar to prokaryotic cell division. Both are double membrane organelles - the inner membrane descended from the ancestral guest cell, and the outer membrane descended from the vacuole membrane that was formed around the guest.

Other organelles may also be the product of endosymbiosis. Some centrioles and basal bodies have naked DNA as part of their structure.

There are many modern examples of endosymbiosis involving organisms that can live together or live independently. The same was probably true of the ancestors of endosymbiotic organelles in the distant past.